



Intravitreal Aflibercept 8 mg in Patients with Polypoidal Choroidal Vasculopathy (PCV): A Phase 3 PULSAR Trial Subgroup Analysis

**Timothy YY Lai,¹ Tien Y Wong,² Jeffrey S Heier,³ Xin Zhang,⁴ Andrea Schulze,⁵
Zoran Hasanbasic,⁴ Sergio Leal,⁴ on behalf of the PULSAR study investigators**

¹The Chinese University of Hong Kong, Hong Kong

²Singapore Eye Research Institute, Singapore National Eye Centre, Singapore; Tsinghua Medicine, Tsinghua University, Beijing, China

³Ophthalmic Consultants of Boston, Boston, MA, USA

⁴Bayer Consumer Care AG, Basel, Switzerland

⁵Bayer AG, Berlin, Germany

Disclosures



- **TYYL:** Research grants (S) from Bayer, Chengdu Kanghong, Novartis, and Roche; consulting fees (C) from Bayer, Boehringer Ingelheim, Iveric Bio, Novartis, Oculis, and Roche; and lecture fees (L) from Bayer, Chengdu Kanghong, Novartis, and Roche
 - **TYW:** Consultant fees from Bayer, Boehringer-Ingelheim, Eden Ophthalmic, Genentech, Iveric Bio, Novartis, Roche, Shanghai Henlius, and Zhaoke Pharmaceutical. **JSH:** Consulting fees from 4DMT, Abpro, Adverum, Affamed, AGTC, Akouos, Allegro, Annexon, Apellis, Asclepix, Bausch & Lomb, Biovisics, Clearside, Curacle, DTx Pharma, Genentech/Roche, Glaukos, Gyroscope, Immunogen, Iveric, Janssen R&D, jCyte, Kriya, Nanoscope, NGM, Notal Vision, Novartis, Ocular Therapeutix, Ocuphire, OcuTerra, Olix, ONL Therapeutics, Palatin, Perceive, Ray Therapeutics, Regeneron, Regenxbio, RetinAI, RevOpsis, Stealth, Thea, Vanotech; research funding from Annexon, Apellis, AsclepiX, Bayer, Genentech/Roche, Gyroscope, Iveric, Kodiak, NGM, Notal Vision, Regeneron, Regenxbio; Equity in Adverum, Aldeyra, Allegro, Aviceda, DTx Pharma, jCyte, Ocuphire, Ocular Therapeutix, RevOpsis, Vinci, and Vitranu; and member of the Board of Directors of Ocular Therapeutix. **XZ, ZH, and SL:** Employees of Bayer Consumer Care AG. **AS:** Employee of Bayer AG
- The PULSAR study (NCT04423718) was sponsored by Bayer AG (Leverkusen, Germany) and co-funded by Regeneron Pharmaceuticals, Inc. (Tarrytown, NY, USA). The sponsor participated in the design and conduct of the study, analysis of the data, and preparation of this presentation
- Study disclosures: This study includes research conducted on human patients. Institutional Review Board approval was obtained prior to study initiation
- The results of the PULSAR polypoidal choroidal vasculopathy subgroup were previously presented at the APAO Congress, February 23–26, 2023; ARVO Annual Meeting, April 23–27, 2023; ASRS Annual Meeting, July 28–August 1, 2023; CCOS Congress, September 6–10, 2023; EURETINA Congress, October 5–8, 2023; JCO Congress, October 6–9, 2023; RetSoc Annual Meeting, October 11–14, 2023; and RANZCO Congress, October 20–23, 2023
- Medical writing support, under the direction of the authors, was provided by ApotheCom and funded by Bayer Consumer Care AG (Basel, Switzerland), in accordance with Good Publication Practice (GPP) standards (*Ann Intern Med* 2022;175:1298–1304)

PULSAR Study Design and Dosing Schedule



96-week, multicenter, randomized, double-masked study in patients with treatment-naïve nAMD
Randomized at baseline 1 (2q8) : 1 (8q12) : 1 (8q16)

Key secondary endpoint:
Proportion of patients without IRF and
SRF in the central subfield

Primary endpoint:
Mean change in BCVA
(non-inferiority)

	Day 1	Wk 4	Wk 8	Wk 12	Wk 16	Wk 20	Wk 24	Wk 28	Wk 32	Wk 36	Wk 40	Wk 44	Wk 48
2q8 (n=336)	X	X	X		X	o	X	o	X	o	X	o	X
8q12 (n=335)	X	X	X		o	X	o	o	X	o	o	X	o
8q16 (n=338)	X	X	X		o	o	X	o	o	o	X	o	o

DRM Criteria for Shortening Dosing Interval

- >5-letter loss in BCVA compared with Week 12, due to persistent or worsening nAMD

AND

- >25 µm increase in CST compared with Week 12, or new-onset foveal neovascularization, or foveal hemorrhage

DRM in Year 1

Intervals can only be
shortened

Multiple opportunities to
shorten interval

Minimum interval for
all patients was **Q8**

Week 16 and 20: Patients on **8q12** and **8q16** meeting DRM criteria shortened to q8

Week 24: Patients on **8q16** meeting DRM criteria shortened to q12

Week 32 and 44 for **8q12** and Week 40 for **8q16**: Treatment interval shortened by 4 weeks for patients meeting DRM criteria


Stippled boxes = initial treatment phase; X = active injection; o = sham injection. Note: Table does not reflect all dosing options once a patient's dosing interval is shortened. 2q8, aflibercept 2 mg every 8 weeks after 3 initial monthly injections; 8q12, aflibercept 8 mg every 12 weeks after 3 initial monthly injections; 8q16, aflibercept 8 mg every 16 weeks after 3 initial monthly injections; BCVA, best-corrected visual acuity; CST, central subfield thickness; DRM, dose regimen modification; IRF, intraretinal fluid; nAMD, neovascular age-related macular degeneration; q8, every 8 weeks; q12, every 12 weeks; SRF, subretinal fluid; Wk, week.

PCV in PULSAR



- PULSAR is a global study conducted across 223 sites in 27 countries
- **ICGA** was optional and **conducted in 292 patients in 12 countries^a**
 - PCV present, n=139; PCV absent, n=153

PCV subgroup by race^b




BL demographics and disease characteristics	PCV subgroup (ICGA-confirmed)				Overall population			
	2q8	8q12	8q16	All 8 mg	2q8	8q12	8q16	All 8 mg
	n=54	n=44	n=41	n=85	n=336	n=335	n=338	n=673
Age, years	72.6 (8.2)	72.2 (8.1)	73.2 (8.7)	72.7 (8.3)	74.2 (8.8)	74.7 (7.9)	74.5(8.5)	74.6 (8.2)
Female, %	31.5	50.0	36.6	43.5	56.0	54.3	53.3	53.8
BCVA, ETDRS letters	57.6 (15.5)	56.3 (13.3)	60.1 (11.5)	58.1 (12.5)	58.9 (14.0)	59.9 (13.4)	60.0 (12.4)	59.9 (12.9)
CST, µm	378 (163)	392 (129)	377 (139)	384 (134)	367 (134)	370 (124)	371 (133)	371 (128)
CNV size, mm ²	5.8 (4.7)	5.1 (3.8)	5.2 (4.5)	5.1 (4.2)	6.4 (5.0)	6.0 (4.8)	6.5 (5.5)	6.3 (5.2)


FAS, data are mean (SD) unless otherwise indicated. ICGA images were graded by the reading center. ^aAustralia (n=15); Austria (n=7); France (n=2); Italy (n=22); Japan (n=70); South Korea (n=25); Latvia (n=2); Mainland China (n=65); Singapore (n=1); Spain (n=3); Switzerland (n=3); USA (n=77). ^bNo patients were reported as Black or African American, or multi-racial.

BL, baseline; CNV, choroidal neovascularization; ETDRS, Early Treatment Diabetic Retinopathy Study; FAS, full analysis set; ICGA, indocyanine green angiography; PCV, polypoidal choroidal vasculopathy; SD, standard deviation.

PCV Subgroup: Absolute and Change from Baseline in BCVA through Week 48 (LOCF)



	Mean $\pm$ SD change from BL to Week 48 (LOCF)	Two-sided 95% CI
2q8	$+9.3 \pm 11.1$	6.2, 12.3
8q12	$+9.5 \pm 13.0$	5.5, 13.4
8q16	$+7.5 \pm 7.6$	5.1, 9.9




	Mean $\pm$ SD change from BL to Week 48 (LOCF)	Two-sided 95% CI
2q8	$+7.5 \pm 12.0$	6.2, 8.8
8q12	$+6.1 \pm 13.2$	4.7, 7.6
8q16	$+5.9 \pm 11.8$	4.6, 7.2

FAS, LOCF (last available observed value prior to ICE will be used to impute missing data; ICE will be handled according to sensitivity estimand strategy for continuous endpoints as described).
 PCV subgroup: 2q8 n=54, 8q12 n=44, 8q16 n=41 (at baseline); overall population: 2q8 n=336, 8q12 n=335, 8q16 n=338 (at baseline).
 CI, confidence interval; ICE, intercurrent event; **LOCF**, last observation carried forward.


PCV Subgroup: Proportions of Patients Maintained on q12 and q16 Dosing Intervals through Week 48




PCV Subgroup



8q12 n=41^b




8q16 n=37^b




All aflibercept 8 mg n=78^b

87% of patients receiving aflibercept 8 mg maintained dosing intervals ≥12 weeks


Overall Population



8q12 n=316^b



8q16 n=312^b



All aflibercept 8 mg n=628^b

83% of patients receiving aflibercept 8 mg maintained dosing intervals ≥12 weeks


Values may not add up to 100% due to rounding.

^aPatients shortened based on DRM assessments at some point through Week 48. ^bPatients completing Week 48.



PCV Subgroup: Anatomic Endpoints

Patients without retinal fluid
in central subfield at Wk 16


PCV Subgroup



Absolute CST




Overall Population




"Without retinal fluid" defined as absence of IRF and SRF in central subfield. For fluid: FAS, LOCF (data post-ICE were censored). For CST: FAS, LOCF.
PCV subgroup: 2q8 n=54, 8q12 n=44, 8q16 n=41 (at baseline); overall population: 2q8 n=335, 8q12 n=333, 8q16 n=334 (at baseline).

PCV Subgroup: Additional Analyses


Patients with number of polypoidal lesions^a



Patients with active polypoidal lesions^{a,c}



Absolute macular volume



Data are for patients with PCV who completed Week 48. Screening (Visit 1) occurred before the baseline visit (Visit 2). ^aFor polypoidal lesion data: 2q8, n=51; All 8 mg, n=78. % calculated based on number of patients with known number of polypoidal lesions. ^bAt Wk 48, number of polypoidal lesions unknown for n=6 and n=9 in 2q8 and All 8 mg groups, respectively. ^c"No" active polypoidal lesions defined as no polypoidal lesions present OR IRF and SRF are "absent" or "questionable". ^dAt Wk 48, n=2 unknown each for 2q8 and All 8 mg groups.

PCV Subgroup: Safety – Week 48

TEAE, %	PCV subgroup				Overall population			
	2q8	8q12	8q16	All 8 mg	2q8	8q12	8q16	All 8 mg
	n=54	n=44	n=41	n=85	n=336	n=335	n=338	n=673
Any ocular TEAE	25.9	38.6	31.7	35.3	38.7	38.5	37.6	38.0
Reduced visual acuity	5.6	4.5	0	2.4	6.0	3.6	5.3	4.5
Cataract	2 cases of cataracts ^a				3.0	3.6	3.6	3.6
Retinal hemorrhage	3.7	6.8	2.4	4.7	4.2	3.3	3.0	3.1
Intraocular pressure increased	1.9	4.5	7.3	5.9	2.1	3.3	2.7	3.0
Vitreous floaters	3.7	0	7.3	3.5	3.3	1.2	3.6	2.4
Subretinal fluid	0	0	0	0	3.3	3.0	1.5	2.2
Intraocular inflammation	1 mild case ^a				0.6	1.2	0.3	0.7
Endophthalmitis	0	0	0	0	0	0	0	0
Occlusive retinal vasculitis	0	0	0	0	0	0	0	0

The safety profile of aflibercept 8 mg was comparable to that of aflibercept 2 mg both in the PCV subgroup and overall population of patients with nAMD

Data are from the SAF. TEAEs are AEs occurring from the first injection to 30 days after the last injection (active or sham); ocular TEAEs are those occurring in the study eye.

^aData presented in this way to avoid unintentional patient unmasking.

AE, adverse event; TEAE, treatment-emergent adverse event; SAF, safety analysis set.

Conclusions:

Aflibercept 8 mg Monotherapy in PCV

Aflibercept 8 mg monotherapy^a showed robust efficacy benefits

Robust improvements in BCVA from baseline to Week 48 were observed in the aflibercept 8q12, 8q16, and 2q8 PCV subgroups (+9.5, +7.5, and +9.3 letters, respectively)

At **Week 16**, **fluid was absent** in 67% and 63% of patients with PCV in the pooled aflibercept 8 mg group and aflibercept 2 mg group, respectively

Aflibercept 8 mg and aflibercept 2 mg **markedly reduced** the proportion of patients with PCV with **active polyps** from baseline (97% and 94%) to Week 48 (21% and 22%), respectively

Extended durability

In total, **87%** of patients with PCV treated with aflibercept 8 mg were maintained on dosing intervals **≥12 weeks**, suggesting extended durability compared with aflibercept 2 mg

Comparable safety profile with aflibercept 2 mg

In the PULSAR study, the **safety profile** of aflibercept 8 mg was **similar** in the PCV subgroup and the overall study population, and the safety profile was comparable to that of aflibercept 2 mg^{1,2}

^aWithout active rescue photodynamic therapy.

1. Heier et al. Ophthalmology 2012;119(12):2537-48; 2. Schmidt-Erfurth et al. Ophthalmology 2014;121(1):193-201.