

Subgroup analyses from the Phase 3 PULSAR trial of aflibercept 8 mg in patients with treatment-naïve neovascular age-related macular degeneration

Keyvan Koushan,¹ Sobha Sivaprasad,² Sergio Leal,³
Tobias Machewitz,⁴ Xin Zhang,³ on behalf of the PULSAR study investigators

¹Department of Ophthalmology and Vision Sciences, University of Toronto, Toronto, Ontario, Canada,
and Toronto Retina Institute, Toronto, Ontario, Canada; ²Moorfields Eye Hospital, London, UK;

³Bayer Consumer Care AG, Basel, Switzerland; ⁴Bayer AG, Berlin, Germany

Disclosures

- Keyvan Koushan receives a research grant from Bayer; and consultant fees from Alcon, Allergan, Bayer, and Novartis
- The PULSAR study was sponsored by Bayer AG (Leverkusen, Germany) and co-funded by Regeneron Pharmaceuticals, Inc. (Tarrytown, NY, USA). The sponsor participated in the design and conduct of the study, analysis of the data, and preparation of this presentation
- Study disclosures: This study includes research conducted on human patients. Institutional Review Board approval was obtained prior to study initiation
- The results from the PULSAR study were previously presented at the ARVO Annual Meeting, April 23–27, 2023
- Medical writing support, under the direction of the authors, was provided by ApotheCom and funded by Bayer Consumer Care AG, Basel, Switzerland, in accordance with Good Publication Practice (GPP) standards (*Ann Intern Med* 2022;175:1298–1304)

PULSAR Study Design

Multicenter, randomized, double-masked study in patients with treatment-naïve nAMD
Randomized at baseline 1 (2q8) : 1 (8q12) : 1 (8q16)

2q8

Aflibercept 2 mg every 8 weeks
after 3 initial monthly injections
n=336

8q12

Aflibercept 8 mg every 12 weeks
after 3 initial monthly injections
n=335

8q16

Aflibercept 8 mg every 16 weeks
after 3 initial monthly injections
n=338

Primary endpoint at Week 48
Mean change in BCVA (non-inferiority)

Key secondary endpoint at Week 16
Proportion of patients without IRF and SRF in the center subfield



End of study at Week 96
with optional 1-year extension through Week 156

Subgroups were determined post hoc and subgroup analyses were exploratory:
For example, baseline BCVA, CST, CNV type and race

BCVA, best corrected visual acuity; CNV, choroidal neovascularization; CST, central subfield thickness; IRF, intraretinal fluid; nAMD, neovascular age-related macular degeneration; SRF, subretinal fluid.

PULSAR: Dosing Schedule and Regimen Modification in Year 1

	Day 1	Wk 4	Wk 8	Wk 12	Wk 16	Wk 20	Wk 24	Wk 28	Wk 32	Wk 36	Wk 40	Wk 44	Wk 48
2q8	X	X	X		X	o	X	o	X	o	X	o	X
8q12	X	X	X		o	X	o	o	X	o	o	X	o
8q16	X	X	X		o	o	X	o	o	o	X	o	o

DRM criteria for shortening dosing interval*

- >5-letter loss in BCVA due to persistent or worsening nAMD

AND

- >25-μm increase in CST, or new-onset foveal neovascularization, or foveal hemorrhage

*All assessments compared to Week 12

DRM in Year 1

Intervals can only be **shortened**

Multiple opportunities to shorten interval

Minimum interval for all patients was **q8**

Week 16 and 20: Patients on **8q12** and **8q16** meeting DRM criteria shortened to q8

Week 24: Patients on **8q16** meeting DRM criteria shortened to q12

Week 32 and 44 for **8q12** and Week 36^a and 40 for **8q16**: Treatment interval shortened by 4 weeks for patients meeting DRM criteria


Stippled boxes = initial treatment phase; X = active injection; o = sham injections. Note: Table does not reflect all dosing options once a patient is shortened.

^aAt Week 36, patients on 8q16 who were previously shortened to q12 could have been shortened to q8.

CST, central subfield thickness; DRM, dose regimen modification; Wk, week.

BCVA by Baseline BCVA Categories (ETDRS Letters): At Week 48

Change from baseline at Week 48^a




FAS, LOCF. Error bars denote SD. Some baseline BCVA plus mean change at Week 48 BCVA values do not add up exactly to the Week 48 absolute BCVA values due to rounding.

^aMean (95% CI) change from baseline to Week 48. **ETDRS**, Early Treatment Diabetic Retinopathy Study; **FAS**, full analysis set; **LOCF**, last observation carried forward.

BCVA by Baseline CST Categories (μm): At Week 48

Change from baseline at Week 48^a




FAS, LOCF. Error bars denote SD. Some baseline BCVA plus mean change at Week 48 BCVA values do not add up exactly to the Week 48 absolute BCVA values due to rounding.

^aMean (95% CI) change from baseline to Week 48.

BCVA by Baseline CNV Type: At Week 48

Change from baseline at Week 48^a




FAS, LOCF. Error bars denote SD. Some baseline BCVA plus mean change at Week 48 BCVA values do not add up exactly to the Week 48 absolute BCVA values due to rounding.

^aMean (95% CI) change from baseline to Week 48.


BCVA by Race: At Week 48

Change from baseline at Week 48^a



FAS, LOCF. Error bars denote SD. Some baseline BCVA plus mean change at Week 48 BCVA values do not add up exactly to the Week 48 absolute BCVA values due to rounding. Data are not reported for Black or African American patients due to small sample size (n=2). ^aMean (95% CI) change from baseline to Week 48.

Proportion of Patients Maintaining 8 mg q12 and q16 Treatment Intervals Up to Week 48: Categorized by Race



FAS, LOCF. Data are not reported for Black or African American patients due to small sample size (n=2).

^aPatients completing Week 48.

Conclusions: Baseline Subgroup Analyses at Week 48

- BCVA outcomes were comparable for aflibercept 8 mg vs aflibercept 2 mg among the evaluable subgroups of baseline characteristics
- Regardless of differences in baseline BCVA and BCVA change at Week 48 in some subgroups, absolute BCVA values at Week 48 were comparable
- The majority (70–80%) of White and Asian patients with nAMD treated with intravitreal aflibercept 8 mg maintained q12 or q16 week treatment intervals.^a These proportions are comparable to those for the overall patient populations in the PULSAR study

^aBlack or African American patients were excluded due to small sample size.