

PULSAR subgroups defined by baseline BCVA, CRT, CNV and race: Similar Week 96 improvements with aflibercept 8 mg q12 weeks or longer vs 2 mg q8 weeks

Nitish Mehta,¹ Richard Gale,² Oliver Zeitz,³ Sobha Sivaprasad,⁴ Sergio Leal,⁵ Tobias Machewitz,⁶ Xin Zhang,⁵ on behalf of the PULSAR study investigators

¹Department of Ophthalmology, NYU Langone Health, New York, New York, USA; ²York and Scarborough Teaching Hospital NHS Foundation Trust, York, UK; ³Department of Ophthalmology, Charité Universitätsmedizin Berlin, Germany; ⁴Moorfields Eye Hospital, London, UK; ⁵Bayer Consumer Care AG, Basel, Switzerland; ⁶Bayer AG, Berlin, Germany

Disclosures

- **NM:** has no financial disclosure
- The PULSAR study (NCT04423718) was sponsored by Bayer AG (Leverkusen, Germany) and co-funded by Regeneron Pharmaceuticals, Inc. (Tarrytown, NY, USA). The sponsor participated in the design and conduct of the study, analysis of the data, and preparation of this presentation
- Study disclosures: This study includes research conducted on human patients. Institutional Review Board approval was obtained prior to study initiation
- The data in this presentation were originally presented at the Association for Research in Vision and Ophthalmology (ARVO) Annual Meeting, Seattle, WA, USA, May 5–9, 2024
- Medical writing support, under the direction of the authors, was provided by ApotheCom and funded by Bayer Consumer Care AG (Basel, Switzerland), in accordance with Good Publication Practice (GPP) guidelines (*Ann Intern Med* 2022;175:1298–1304)

PULSAR: Multicenter, randomized, double-masked study

Patients with treatment-naïve nAMD, randomized at baseline

2q8

Aflibercept 2 mg every 8 weeks
after 3 initial monthly injections
n=336

8q12

Aflibercept 8 mg every 12 weeks
after 3 initial monthly injections
n=335

8q16

Aflibercept 8 mg every 16 weeks
after 3 initial monthly injections
n=338

	YEAR 1													YEAR 2											
	Day 1	W4	W8	W12	W16	W20	W24	W28	W32	W36	W40	W44	W48	W52	W56	W60	W64	W68	W72	W76	W80	W84	W88	W92	W96
2q8	X	X	X		X	o	X	o	X	o	X	o	X	o	X	o	X	o	X	o	X	o	X	o	–
8q12	X	X	X		<u>o^a</u>	<u>X^a</u>	o	o	<u>X^a</u>	o	o	<u>X^a</u>	o	o	<u>X^{a,b}</u>	o	o	<u>X^{a,b}</u>	o	o	<u>X^{a,b}</u>	o	o	<u>X^{a,b}</u>	–
8q16	X	X	X		<u>o^a</u>	<u>o^a</u>	<u>X^a</u>	o	o	o	<u>X^a</u>	o	o	o	<u>X^{a,b}</u>	o	o	o	<u>X^{a,b}</u>	o	o	o	<u>X^{a,b}</u>	o	–

Primary endpoint at W48:
Mean change in BCVA
(non-inferiority)

End of study at W96
with optional ~1-year
extension through W156

^aDRM: Interval Shortening During Years 1 and 2

Criteria for interval shortening

- >5-letter loss in BCVA compared with Week 12 due to persistent or worsening nAMD **AND**
- >25 µm increase in CRT compared with Week 12, **OR** new foveal neovascularization, **OR** new foveal hemorrhage

- Patients who met DRM criteria had dosing intervals shortened to q8 at Weeks 16 and 20 or by 4-week increments from Week 24
 - The minimum assigned dosing interval was q8

^bDRM: Interval Extension During Year 2

Criteria for interval extension


- <5-letter loss in BCVA compared with Week 12 **AND**
- No fluid at the central subfield on OCT **AND**
- No new foveal hemorrhage or foveal neovascularization

- Patients who met DRM criteria from Weeks 52 through 96 had dosing intervals extended by 4-week increments
 - The maximum assigned dosing interval was q24

Figure does not reflect all dosing options once a patient's dosing interval is shortened or extended. Stippled boxes = initial treatment phase; X = active injection; o = sham injections. q8, every 8 weeks; q24, every 24 weeks; BCVA, best-corrected visual acuity; CRT, central retinal thickness; DRM, dose regimen modification; nAMD, neovascular age-related macular degeneration; OCT, optical coherence tomography; W, week.

BCVA and CRT through Week 96: Comparable outcomes with aflibercept 8 mg and 2 mg overall


BCVA change from BL^a



LS means^{a,b}
Week 48 Week 96

2q8	+7.0	+6.6
8q12	+6.1	+5.6
8q16	+5.9	+5.5

Absolute CRT^c



LS means^{a,b}
Week 48 Week 96

2q8	-136	-147
8q12	-147	-152
8q16	-147	-149

FAS: 2q8 n=336; 8q12 n=335; 8q16 n=338 (at BL). ^aLS mean values (data post-ICE were censored); ^bLS means were generated using MMRM, with baseline BCVA measurement (for BCVA analysis) or BL CRT measurement (for CRT analysis) as a covariate, and treatment group (aflibercept 2q8, 8q12, 8q16), visit, and stratification variables (geographic region [Japan vs. Rest of World] and BL BCVA [<60 vs. ≥ 60]) as fixed factors, and interaction terms for BL and visit and for treatment and visit. ^cObserved values.

BL, baseline; ETDRS, Early Treatment of Diabetic Retinopathy Study; FAS, full analysis set, ICE, intercurrent event; LS, least squares; MMRM, mixed model for repeated measures.

Baseline characteristics

	2q8	8q12	8q16	All 8mg	Total
Randomized, n	336	335	338	673	1009
Age, years	74.2 (8.8)	74.7 (7.9)	74.5 (8.5)	74.6 (8.2)	74.5 (8.4)
Female, %	56.0	54.3	53.3	53.8	54.5
Race, %					
Asian	24.7	22.1	22.8	22.4	23.2
Black or African American	0.6	0.6	0	0.3	0.4
White	74.1	76.4	76.9	76.7	75.8
Not reported	0.6	0.6	0.3	0.4	0.5
BCVA, ETDRS letters	58.9 (14.0)	59.9 (13.4)	60.0 (12.4)	59.9 (12.9)	59.6 (13.3)
CRT, μm	367 (134)	370 (124)	371 (133)	371 (128)	370 (130)
CNV size, mm^2	6.4 (5.0)	6.0 (4.8)	6.5 (5.5)	6.3 (5.2)	6.3 (5.1)
CNV type, %					
Minimally classic CNV	18.2	16.7	20.1	18.4	18.3
Occult-only CNV	57.1	58.8	55.0	56.9	57.0
Predominantly classic CNV	21.1	21.2	19.8	20.5	20.7

BCVA outcomes with aflibercept 8 mg and 2 mg at Week 96 are independent of BL BCVA

Mean (95% CI) BCVA (ETDRS letters) change from BL at Week 96

BL BCVA ≤54


+11.7 (8.8, 14.7)
+10.4 (7.0, 13.7)
+6.5 (2.9, 10.0)

BL BCVA 55–73

+5.9 (4.2, 7.6)
+4.1 (2.1, 6.1)
+5.9 (4.4, 7.5)

BL BCVA ≥74


+1.5 (-0.8, 3.9)
+0.9 (-2.6, 4.3)
+1.1 (-2.1, 4.4)



FAS, LOCF. N values for BL. Analyses were not adjusted for multiplicity or to allow for differences in BL BCVA. CI, confidence interval; LOCF, last observation carried forward.

CRT improvements with aflibercept 8 mg and 2 mg at Week 96 are independent of BL BCVA


	BL BCVA ≤54	BL BCVA 55–73	BL BCVA ≥74
Mean CRT (μm) change (95% CI) from BL at Week 96	-224 (-255, -193)	-115 (-129, -100)	-58 (-75, -42)
	-254 (-283, -224)	-109 (-121, -96)	-76 (-101, -51)
	-237 (-270, -205)	-112 (-126, -97)	-82 (-107, -58)




Last assigned treatment intervals at Week 96 categorized by BL BCVA

BL BCVA categories


≤54 ETDRS letters
(n=166)



55–73 ETDRS letters
(n=338)



≥74 ETDRS letters
(n=79)



All patients receiving aflibercept 8 mg

Last assigned treatment interval:

■ q8

■ q12

■ q16

■ q20

■ q24^a


Data shown for patients who completed 96 weeks of treatment.

Values may not add up to 100% due to rounding.

^aPatients assigned to a 24-week dosing interval did not have enough time to complete the interval within the 96-week study period.


BCVA outcomes with aflibercept 8 mg and 2 mg at Week 96 are independent of BL CRT

	≤278 μm	279–343 μm	344–422 μm	≥423 μm
Mean (95% CI) BCVA (ETDRS letters) change from BL at Week 96	+4.8 (2.2, 7.4)	+6.3 (3.8, 8.8)	+6.9 (4.1, 9.7)	+10.4 (7.2, 13.6)
	+3.1 (0.7, 5.4)	+4.2 (0.3, 8.0)	+5.2 (2.4, 8.1)	+9.3 (5.7, 12.9)
	+7.4 (5.3, 9.5)	+5.0 (2.6, 7.4)	+4.3 (0.9, 7.7)	+4.8 (1.2, 8.3)




CRT improvements with aflibercept 8 mg and 2 mg at Week 96 are independent of BL CRT

	≤278 μm	279–343 μm	344–422 μm	≥423 μm
Mean (95% CI) CRT (μm) change from BL at Week 96	-32 (-45, -19)	-79 (-91, -66)	-146 (-161, -132)	-310 (-335, -285)
	-49 (-58, -40)	-80 (-94, -67)	-143 (-157, -128)	-308 (-335, -280)
	-41 (-50, -33)	-82 (-94, -69)	-142 (-156, -128)	-317 (-347, -288)



Last assigned treatment intervals at Week 96 categorized by BL CRT

BL CRT categories




Data shown for patients who completed 96 weeks of treatment. Values may not add up to 100% due to rounding.

^aPatients assigned to a 24-week dosing interval did not have enough time to complete the interval within the 96-week study period.

BCVA outcomes with aflibercept 8 mg and 2 mg at Week 96 are independent of BL CNV size

Mean (95% CI) BCVA (ETDRS letters) change from BL at Week 96


	<1.3 mm ²	1.3–<4 mm ²	4–<5 mm ²	≥5 mm ²
2q8	+10.7 (6.3, 15.2)	+8.2 (5.8, 10.6)	+5.5 (1.2, 9.8)	+6.0 (3.9, 8.1)
8q12	+7.0 (2.8, 11.2)	+7.9 (5.1, 10.7)	+5.5 (1.1, 9.8)	+3.7 (1.2, 6.3)
8q16	+7.8 (3.9, 11.7)	+8.2 (5.9, 10.5)	+4.2 (1.1, 7.3)	+3.5 (1.2, 5.8)



CRT improvements with aflibercept 8 mg and 2 mg at Week 96 are independent of BL CNV size


Mean (95% CI)
CRT (μm)
change from
BL at
Week 96

<1.3 mm ²	1.3–<4 mm ²	4–<5 mm ²	≥ 5 mm ²
-119 (-158, -80)	-138 (-164, -111)	-157 (-205, -109)	-144 (-164, -123)
-102 (-133, -71)	-155 (-184, -126)	-135 (-173, -97)	-157 (-177, -137)
-126 (-165, -86)	-135 (-164, -106)	-109 (-144, -75)	-165 (-186, -144)



Last assigned treatment intervals at Week 96 categorized by BL CNV size

Baseline CNV size





Data shown for patients who completed 96 weeks of treatment. Values may not add up to 100% due to rounding.

^aPatients assigned to a 24-week dosing interval did not have enough time to complete the interval within the 96-week study period.

Visual and anatomic outcomes with aflibercept 8 mg and 2 mg at Week 96 are independent of race

Mean (95% CI) BCVA (ETDRS letters) and CRT (μm) change from BL at Week 96

Asian		White		Asian		White	
+7.5	(4.8, 10.3)	+6.9	(5.3, 8.6)	-144	(-176, -111)	-138	(-154, -123)
+8.9	(5.1, 12.8)	+4.4	(2.6, 6.1)	-147	(-179, -115)	-147	(-163, -132)
+7.2	(4.8, 9.6)	+4.8	(3.1, 6.5)	-140	(-169, -112)	-146	(-163, -129)




	2q8	8q12	8q16	2q8	8q12	8q16
BL	59.2	57.7	58.1	59.0	60.3	60.7
Week 48	66.7	67.0	66.8	66.5	65.5	65.7


Last assigned treatment intervals at Week 96 categorized by race

Race

Asian
(n=130)



White
(n=447)



All patients receiving aflibercept 8 mg

Last assigned treatment interval:



Data shown for patients who completed 96 weeks of treatment. Values may not add up to 100% due to rounding.


Data are not reported for Black or African American patients due to small sample size (n=2).

^aPatients assigned to a 24-week dosing interval did not have enough time to complete the interval within the 96-week study period.


Conclusions

- At Week 96, mean absolute and change in BCVA and CRT values were similar after treatment with aflibercept 8 mg with extended treatment intervals compared with 2 mg every 8 weeks **across all subgroups based on** baseline BCVA, CRT, CNV size, and race.
- Proportion of patients with maintained, extended, or shortened **treatment intervals** through Week 96 was **also independent of baseline characteristics**

Week 96 BCVA outcomes according to BL BCVA



Last assigned treatment interval according to BL BCVA^a



^aValues may not add up to 100% due to rounding. ^bPatients assigned to a 24-week dosing interval did not have enough time to complete the interval within the 96-week study period.