



Efficacy and safety outcomes from the FIREFLEYE next study of children 3 years of age with retinopathy of prematurity treated with intravitreal aflibercept versus laser in the randomized FIREFLEYE study

Andreas Stahl,¹ Hidehiko Nakanishi,² Domenico Lepore,³ Wei-Chi Wu,⁴ Noriyuki Azuma,⁵ Carlos Jacas,⁶ Aditya Athanikar,⁷ Robert Vitti,⁷ Karen Chu,⁷ Pablo Iveli,⁸ Fei Zhao,⁹ Sarah Schlieff,¹⁰ Sergio Leal,¹¹ Katja Brandau,⁸ Thomas Miller,¹² Evra Köföncü,¹⁰ Alistair Fielder,¹³ on behalf of the FIREFLEYE next investigators

¹Department of Ophthalmology, University Medicine Greifswald, Greifswald, Germany; ²Research and Development Center for New Medical Frontiers, Department of Advanced Medicine, Division of Neonatal Intensive Care Medicine, Kitasato University School of Medicine, Sagami-hara, Japan; ³Department of Neuroscience, Sensory Organs and Thorax, Catholic University of the Sacred Heart, A. Gemelli Foundation IRCCS, Rome, Italy; ⁴Department of Ophthalmology, Linkou Chang Gung Memorial Hospital, and College of Medicine, Chang Gung University, Taoyuan, Taiwan; ⁵Department of Developmental and Regenerative Biology, Medical Research Institute, Tokyo Medical and Dental University, Tokyo, Japan; ⁶Department of Psychiatry, Hospital Universitari Vall d'Hebron, Vall d'Hebron Barcelona Hospital Campus, Barcelona, Spain; ⁷Regeneron Pharmaceuticals Inc., Tarrytown, NY, USA; ⁸Bayer AG, Wuppertal, Germany; ⁹Bayer Inc., Mississauga, ON, Canada; ¹⁰Bayer AG, Berlin, Germany; ¹¹Bayer Consumer Care AG, Basel, Switzerland; ¹²Bayer Healthcare, LLC, Whippany, NJ, USA; ¹³Department of Optometry and Visual Science, City University London, London, UK

Presented at the 24th European Society of Retina Specialists (EURETINA), Barcelona, Spain, September 19–22, 2024


Disclosures

- **Andreas Stahl:** Consultant for Allergan, Apellis, Bayer, Novartis, and Roche
 - **HN, NA, and CJ:** Expenses from Bayer. **DL and AF:** Consultants for Bayer and Novartis. **W-CW:** Consultant for Allergan. **AA, RV, and KC:** Employees of Regeneron. **PI, FZ, SS, SL, KB, TM, and EK:** Employees of Bayer
- The ongoing FIREFLEYE next study is sponsored by Bayer AG (Leverkusen, Germany). The sponsor participated in the design and conduct of the study, analysis of the data, and preparation of this abstract
- This study includes research conducted on human patients. Institutional Review Board approval was obtained prior to study initiation
- Medical writing support, under the direction of the author, was provided by ApotheCom and funded by Bayer Consumer Care AG, Basel, Switzerland, in accordance with Good Publication Practice (GPP) guidelines (*Ann Intern Med* 2022;175:1298–1304)



FIREFLEYE next (NCT04015180) study design

FIREFLEYE next is the first multinational, ongoing, Phase 3b study assessing ocular and further clinical outcomes, including growth and neurodevelopmental outcomes, **through 5 years of age** following treatment of acute-phase ROP with intravitreal aflibercept 0.4 mg vs. laser photocoagulation in the **24-week, Phase 3 FIREFLEYE study**¹



Endpoints included in pre-planned interim analysis (3 years of chronological age)

- Absence of **active ROP**
- Absence of **unfavorable structural outcomes**
- **Completion of retinal vascularization**
- **Treatment need** for ROP complications
- **Refractive spherical equivalent**
- **BCVA function** (Snellen equivalent)
- Ability to **fix and follow a 5-cm toy**
- Outcomes of **growth and neurodevelopmental tests** (optional [WPPSI-IV, VABS-II])

^aThree study treatment-unrelated deaths in FIREFLEYE and the parents/carers of 10 children (aflibercept 0.4 mg, 6; laser, 4) did not consent to enrollment in FIREFLEYE next. **BCVA**, best corrected visual acuity; **n**, number; **ROP**, retinopathy of prematurity; **VABS-II**, Vineland Adaptive Behavior Scales, Second Edition; **WPPSI-IV**, Wechsler Preschool and Primary Scale of Intelligence, Fourth Edition. 1. Stahl A, et al. JAMA 2022;328:348–59. 3



Baseline characteristics

| | Aflibercept 0.4 mg (n=66) | Laser (n=34) | Total (N=100) |
|---|------------------------------|-----------------|------------------|
| Male, % | 54.5 | 50.0 | 53.0 |
| Gestational age, w and d | 26w 4d ±2.1 | 26w 0d ±1.7 | 26w 3d ±1.9 |
| Gestational age group, % | | | |
| <24 weeks | 4.5 | 8.8 | 6.0 |
| ≥24 to <27 weeks | 59.1 | 64.7 | 61.0 |
| ≥27 weeks | 36.4 | 26.5 | 33.0 |
| Birth weight, g | 882.2 ±286.9 | 819.5 ±238.6 | 860.9 ±271.9 |
| Body weight at baseline treatment in FIREFLEYE, g | 2045.8 ±675.8 | 1843.8 ±569.2 | 1977.1 ±645.8 |
| Chronological age at FIREFLEYE next entry, months | 9.0 ±1.6 | 9.1 ±1.7 | 9.0 ±1.6 |



84% of children had **no ROP** at FIREFLEYE next study entry

| Detailed ROP classification by investigator, % | At FIREFLEYE study entry | | | At FIREFLEYE next study entry | | |
|--|------------------------------|-----------------|------------------|-------------------------------|-----------------|------------------|
| | Aflibercept 0.4 mg (n=75) | Laser (n=38) | Total (N=113) | Aflibercept 0.4 mg (n=66) | Laser (n=34) | Total (N=100) |
| Absence of ROP | 0 | 0 | 0 | 83.3 | 85.3 | 84.0 |
| Zone I (excluding AP-ROP) | 20.0 | 18.4 | 19.5 | 4.5 | 8.8 | 6.0 |
| Stage 1 | 1.3 | 0 | 0.9 | 1.5 | 2.9 | 2.0 |
| Stage 2 | 2.7 | 5.3 | 3.5 | 0 | 0 | 0 |
| Stage 3 | 4.0 | 2.6 | 3.5 | 0 | 0 | 0 |
| Stage 3+ | 12.0 | 10.5 | 11.5 | 0 | 0 | 0 |
| Stage 4A | 0 | 0 | 0 | 1.5 | 0 | 1.0 |
| Stage 4B | 0 | 0 | 0 | 1.5 | 2.9 | 2.0 |
| Missing | 0 | 0 | 0 | 0 | 2.9 | 1.0 |
| Zone II (excluding AP-ROP) | 61.3 | 68.4 | 63.7 | 6.1 | 5.9 | 6.0 |
| Stage 1 | 0 | 0 | 0 | 1.5 | 2.9 | 2.0 |
| Stage 2 | 0 | 2.6 | 0.9 | 3.0 | 0 | 2.0 |
| Stage 2+ | 9.3 | 13.2 | 10.6 | 0 | 0 | 0 |
| Stage 3+ | 52.0 | 52.6 | 52.2 | 0 | 0 | 0 |
| Missing | 0 | 0 | 0 | 1.5 | 2.9 | 2.0 |
| Zone III (excluding AP-ROP) | 0 | 0 | 0 | 6.1 | 0 | 4.0 |
| Stage 1 | 0 | 0 | 0 | 4.5 | 0 | 3.0 |
| Missing | 0 | 0 | 0 | 1.5 | 0 | 1.0 |
| AP-ROP | 18.7 | 13.2 | 16.8 | 0 | 0 | 0 |
| Zone I | 16.0 | 10.5 | 14.2 | 0 | 0 | 0 |
| Zone II | 2.7 | 2.6 | 2.7 | 0 | 0 | 0 |



Unfavorable structural outcomes, ROP recurrence, and treatment for ROP complications in FIREFLEYE next

| Unfavorable structural outcomes | Aflibercept 0.4 mg (n=66) | | Laser (n=34) | |
|--|--|--|--|--|
| | At any time until 2 years of chronological age | At any time until 3 years of chronological age | At any time until 2 years of chronological age | At any time until 3 years of chronological age |
| Number of children, % | 100.0 | 100.0 | 100.0 | 100.0 |
| None | 93.9 | 93.9 | 94.1 | 94.1 |
| Retinal detachment | 4.5 | 6.1 | 2.9 | 2.9 |
| Macular dragging | 1.5 | 1.5 | 2.9 | 2.9 |
| Macular fold | 1.5 | 1.5 | 0 | 0 |
| Retrolental opacity | 1.5 | 1.5 | 0 | 0 |
| Any unfavorable structural outcome | 6.1 | 6.1 | 5.9 | 5.9 |
| Number of treated eyes, % | 100.0 | 100.0 | 100.0 | 100.0 |
| None | 94.5 | 94.5 | 95.3 | 95.3 |
| Retinal detachment | 3.9 | 4.7 | 1.6 | 1.6 |
| Macular dragging | 1.6 | 1.6 | 3.1 | 3.1 |
| Macular fold | 1.6 | 1.6 | 0 | 0 |
| Retrolental opacity | 1.6 | 1.6 | 0 | 0 |
| Any unfavorable structural outcome | 5.5 | 5.5 | 4.7 | 4.7 |
| Recurrence of ROP after entry into FIREFLEYE next^a | Between entry and 2 years of age | Between entry and 3 years of age | Between entry and 2 years of age | Between entry and 3 years of age |
| n | 64 | 60 | 32 | 30 |
| Recurrence, % | 1.5 | 1.7 | 0 | 0 |

- **No disease reactivation occurred after 50 weeks of chronological age**
- In total, **4 patients were treated** after entry into FIREFLEYE next for ROP complications, all before 1 year of age (including 2 patients with pre-existing bilateral retinal detachment, 1 with reactivated plus disease^b, and 1 with retinal neovascularization not further specified^c)
 - 1 patient showed retinal detachment at age 3 (in the progression of macular fold reported at ages 1 and 2 years)


^aPost-hoc analysis. A child was considered as having ROP recurrence if: the inclusion criteria of FIREFLEYE (or worse) were reported and a previous assessment (either in FIREFLEYE or in FIREFLEYE next) of ROP not requiring treatment (according to the inclusion criteria) was available. ^bZone I, both eyes, treated at around 43 weeks of age. ^cTreated around 50 weeks of chronological age.



Ophthalmic outcomes at 3 years of age

| No./Total No. (%) | Aflibercept 0.4 mg | Laser |
|---|--------------------|--------------|
| Ocular findings, eyes | | |
| Nystagmus | 4/116 (3.4) | 2/58 (3.4) |
| Absence of manifest strabismus | 98/116 (84.5) | 50/58 (86.2) |
| Amblyopia | 1/116 (0.9) | 2/58 (3.4) |
| Cataract | 1/116 (0.9) | 0 |
| Optic nerve atrophy | 2/116 (1.7) | 0 |
| Ability to fix and follow a 5-cm toy | 112/116 (96.6) | 57/58 (98.3) |
| Eyes with complete retinal vascularization^a | | |
| At 1 year of chronological age | 89/128 (69.5) | - |
| At 2 years of chronological age | 97/121 (80.2) | - |
| At 3 years of chronological age | 89/111 (80.2) | - |
| BCVA (Snellen equivalent score), patients | | |
| ≥20/200 | 44/45 (97.8) | 23/23 (100) |
| ≥20/40 | 30/45 (66.7) | 11/23 (47.8) |

Retinal vascularization after aflibercept treatment appeared to be complete in 80% of eyes by 2 years of age



Fix and follow a 5 cm toy

Aflibercept 0.4 mg, **97%**; laser, **98%**

BCVA ≥20/40

Aflibercept 0.4 mg, **67%**; laser, **48%**

Myopia

Mild in both groups and **less pronounced with aflibercept**

Aflibercept 0.4 mg, **-0.4 (3.1) D**;
Laser **-2.2 (4.2) D**

High myopia


Aflibercept 0.4 mg **10 eyes (9%)**;
Laser **14 eyes (24%)**

Very high myopia

Aflibercept 0.4 mg **1 eye (1%)**;
Laser **10 eyes (17%)**



Growth parameters up to 3 years of age



No clinically relevant differences in growth parameters were observed between both groups through 3 years of chronological age, and **results are in line with what is expected in this prematurely born pediatric population**



Adverse events (during FIREFLEYE next through 3 years of age)

| | Aflibercept 0.4 mg (n=66) | Laser (n=34) |
|---|---------------------------|--------------------|
| Any AE, n (%) | 59 (89.4) | 29 (85.3) |
| Ocular AEs | 33 (50.0) | 11 (32.4) |
| Ocular AEs in eyes formerly treated in FIREFLEYE | 32 (48.5) | 10 (29.4) |
| Systemic AEs | 56 (84.8) | 29 (85.3) |
| AEs related to aflibercept 0.4 mg | 2 (3.0) | 1 (2.9) |
| AEs related to laser treatment | 3 (4.5) | 6 (17.6) |
| Ocular AEs in eyes formerly treated in FIREFLEYE (≥10% occurrence in any group), n (%) | | |
| Astigmatism | 13 (19.7) | 5 (14.7) |
| Myopia | 9 (13.6) | 5 (14.7) |
| Strabismus ^a | 9 (13.6) | 2 (5.9) |
| Any SAE, n (%) | 21 (31.8) | 14 (41.2) |
| Ocular SAEs in eyes formerly treated in FIREFLEYE | 6 (9.1) | 0 |
| Systemic SAEs | 19 (28.8) | 14 (41.2) |
| SAEs related to aflibercept 0.4 mg | 1 (1.5) | 0 |
| Death | 0 | 0 |
| Ocular SAEs, n (%) | 6 (9.1) | 0 |
| Optic atrophy | 2 (3.0) | 0 |
| Retinal detachment | 2 (3.0) | 0 |
| Retinal neovascularization | 2 (3.0) | 0 |
| Retinopathy of prematurity | 1 (1.5) | 0 |
| Vitreous opacities | 1 (1.5) | 0 |
| Retinoblastoma | 1 (1.5) | 0 |
| Systemic SAEs (≥5% occurrence in any arm), n (%) | | |
| Cerebral palsy | 2 (3.0) | 4 (11.8) |
| Bronchiolitis | 2 (3.0) | 2 (5.9) |
| Bronchospasm | 0 | 2 (5.9) |



Ocular and systemic AEs were consistent with those expected in children born preterm and who developed severe ROP, and no new safety concerns were identified

^aReported as an adverse event. No imbalance of “clinically manifest strabismus” (reported as an efficacy parameter) between groups.
AE, adverse event; SAE, serious adverse event.



Conclusions

FIREFLEYE next is the **first prospective, controlled, Phase 3b study** evaluating **long-term efficacy and safety outcomes** after treatment of **acute-phase ROP with aflibercept 0.4 mg versus laser photocoagulation** (final results through 5 years of age expected for 2026)

Through 3 years of chronological age, **efficacy outcomes were well sustained, and no ocular or systemic safety concerns, including outcomes of growth, were identified:**

Efficacy

- **Disease reactivation** after aflibercept 0.4 mg **was rare**
 - No disease reactivation occurred after 50 weeks of chronological age. One patient showed retinal detachment at age 3 years in the progression of macular fold reported at ages 1 and 2 years of chronological age
- No patient received treatment in the post-acute phase later than 50 weeks of chronological age
- **Retinal vascularization** after aflibercept 0.4 mg **appeared to be complete in 80% of eyes by 2 years** of chronological age
- **Visual function was age-appropriate**, and **myopia** was rarer and **less severe in the aflibercept 0.4 mg group** than the laser group

Safety

- **No ROP treatment-specific effects** on **growth** outcomes through 3 years of age
- No ocular or systemic safety concerns through 3 years of age were identified



Overall, aflibercept 0.4 mg injection therapy in very/extremely preterm or very low birthweight patients with acute-phase ROP (as approved^a) was **effective and generally well tolerated through 3 years of age**

^aAflibercept has been approved for treatment of ROP in Japan (September 2022),¹ the European Union (December 2022),² Switzerland,² Great Britain,² the USA (February 2023),³ and Brazil (April 2023).⁴

1. Bayer, 2023. Available at: <https://www.bayer.com/media/en-us/eylea-approved-in-japan-for-treatment-of-preterm-infants-with-retinopathy-of-prematurity/> [Accessed July 2024]; 2. Bayer AG, 2023. Available at: https://www.ema.europa.eu/en/documents/product-information/eylea-epar-product-information_en.pdf [Accessed July 2024]; 3. Regeneron Pharmaceuticals, 2023. Available at: https://www.regeneron.com/downloads/eylea_fpi.pdf [Accessed July 2024]; 4. Anvisa NHTSA, 2023. Available at: <https://www.gov.br/anvisa/pt-br/assuntos/medicamentos/novos-medicamentos-e-indicacoes/eylea-aflibercepte-nova-indicacao> [Accessed July 2024].



Thank you to all FIREFLEYE next patients, parents, caregivers, and investigators

Final results through 5 years of age are expected in 2026

For more information, please contact:

Andreas Stahl

andreas.stahl@med.uni-greifswald.de

