



photon

Aflibercept 8 mg for Diabetic Macular Edema: 96-Week Results From the Phase 2/3 PHOTON Trial

Diana V. Do,¹ on behalf of the PHOTON study investigators

¹Byers Eye Institute, Stanford University School of Medicine, Palo Alto, CA, USA

Disclosures

- Diana V. Do is a consultant to Boehringer Ingelheim, Genentech, Kodiak Sciences, Kriya, and Regeneron Pharmaceuticals, Inc.; has received research funding from Boehringer Ingelheim, Genentech, Kriya, and Regeneron Pharmaceuticals, Inc.; and has stock options from Kodiak Sciences
- This study was sponsored by Regeneron Pharmaceuticals, Inc. (Tarrytown, NY, USA) and co-funded by Bayer AG (Leverkusen, Germany). The sponsors participated in the design and conduct of the study, analysis of the data, and preparation of this presentation
- Study disclosures: This study includes research conducted on human patients. Institutional Review Board approval was obtained prior to study initiation

PHOTON Study Design

photon

DME

Multi-center, randomized, double-masked study in adult patients with center-involved DME^a

Randomized 1 (2q8) : 2 (8q12) : 1 (8q16)

Note: 2 mg arm received 5 initial monthly injections versus 8 mg arms, which received only 3 initial monthly injections

2q8

Aflibercept 2 mg every 8 weeks
after 5 initial monthly injections
n=167

8q12

8 mg every 12 weeks after
3 initial monthly injections
n=328

8q16

8 mg every 16 weeks after
3 initial monthly injections
n=163

Primary endpoint at Week 48
Mean change in BCVA (non-inferiority)

End of study at Week 96
with optional 1-year extension through Week 156

^aTreatment-naïve and previously treated patients aged ≥18 years with type 1 or type 2 diabetes, DME with central involvement with CRT ≥300 µm in the study eye, and BCVA of 78-24 letters (Snellen equivalent of 20/32-20/320) with decreased vision due to DME.
BCVA, best-corrected visual acuity; CRT, central retinal thickness; DME, diabetic macular edema.

PHOTON: Dosing Schedule and Dose Regimen Modification



DME
Primary
Endpoint

YEAR 1	Day 1	Week 4	Week 8	Week 12	Week 16	Week 20	Week 24	Week 28	Week 32	Week 36	Week 40	Week 44	Week 48
2q8	X	X	X	X	X	o	X	o	X	o	X	o	X
8q12	X	X	X	o	o ^a	X ^a	o	o	X ^a	o	o	X ^a	o
8q16	X	X	X	o	o ^a	o ^a	X ^a	o	o	o	X ^a	o	o

YEAR 2	Week 52	Week 56	Week 60	Week 64	Week 68	Week 72	Week 76	Week 80	Week 84	Week 88	Week 92	Week 96
2q8	o	X	o	X	o	X	o	X	o	X	o	o
8q12	o	X ^{a, b}	o	o	X ^{a, b}	o	o	X ^{a, b}	o	o	X ^{a, b}	o
8q16	o	X ^{a, b}	o	o	o	X ^{a, b}	o	o	o	X ^{a, b}	o	o

^aDRM: Interval Shortening During Years 1 and 2

- Criteria for interval shortening:
 - >10-letter loss in BCVA from Week 12 due to persistent or worsening DME **AND**
 - >50-μm increase in CRT from Week 12
- Patients who met DRM criteria had dosing intervals shortened to Q8 at **Weeks 16 and 20** or by 4-week increments from **Week 24**
 - The minimum interval was Q8

^bDRM: Interval Extension During Year 2

- Criteria for interval extension:
 - <5-letter loss in BCVA from Week 12 **AND**
 - CRT <300 μm (or <320 μm on Spectralis)
- Patients who met DRM criteria beginning at **Week 52** had dosing intervals extended by 4-week increments
 - The maximum assigned interval was Q24

Figure does not reflect all dosing options once a patient's interval is shortened or extended. Stippled boxes = initial treatment phase; X = active injection; o = sham injection. DRM, dose regimen modification.

Patient Disposition

	2q8	8q12	8q16	Total
N (FAS/SAF)	167	328	163	658
Completion rate at Week 48 (%)	94.0%	91.2%	95.1%	92.9%
Completion rate at Week 96 (%)	83.2%	77.8%	84.8%	80.9%
Reasons for discontinuation through Week 96 (%)				
Consent withdrawal	5.4%	5.2%	4.9%	5.2%
Death	5.4%	5.5%	3.0%	4.8%
Lost to follow-up	3.0%	5.8%	4.3%	4.7%
Decision by the investigator	1.2%	2.7%	1.8%	2.1%
Adverse event	0.6%	2.7%	1.2%	1.8%
Noncompliance with protocol	1.2%	0.3%	0	0.5%

Data are mean (SD) unless otherwise indicated.

BMI, body mass index; FAS, full analysis set; SAF, safety analysis set; SD, standard deviation.

Patient Baseline Characteristics

	2q8	8q12	8q16	Total	DME
N (FAS/SAF)	167	328	163	658	
Age (years)	63.0 (9.8)	62.1 (11.1)	61.9 (9.5)	62.3 (10.4)	
Female (%)	44.9%	36.0%	39.3%	39.1%	
Race (%)					
White	67.1%	70.4%	78.5%	71.6%	
Black or African American	10.8%	10.7%	5.5%	9.4%	
Asian	18.0%	14.6%	14.1%	15.3%	
Other	2.4%	3.0%	0.6%	2.4%	
Not reported	1.8%	1.2%	1.2%	1.4%	
Hispanic or Latino (%)	18.6%	16.5%	20.9%	18.1%	
Duration of diabetes (years)	15.9 (10.0)	15.1 (10.0)	15.7 (10.7)	15.5 (10.2)	
Hemoglobin A1c (%)	8.1 (1.5)	7.9 (1.5)	7.8 (1.5)	8.0 (1.5)	
History of hypertension (%)	77.8%	77.4%	79.8%	78.1%	
BMI (kg/m ²)	29.9 (6.5)	30.4 (6.2)	31.0 (6.1)	30.5 (6.2)	

Data are mean (SD) unless otherwise indicated.

BMI, body mass index; FAS, full analysis set; SAF, safety analysis set; SD, standard deviation.


Baseline Characteristics of the Study Eye

	2q8	8q12	8q16	Total
N (FAS/SAF)	167	328	163	658
BCVA (ETDRS letters)	61.5 (11.2)	63.6 (10.1)	61.4 (11.8)	62.5 (10.9)
Snellen equivalent	20/63	20/50	20/63	20/63
20/32 (>73 to 78 letters)	12.0%	18.0%	14.1%	15.5%
20/40 or worse (≤73 letters)	88.0%	82.0%	85.9%	84.5%
CRT (μm)	457.2 (144.0)	449.1 (127.4)	460.3 (117.8)	454.0 (129.5)
Prior treatment for DME (%)	44.3%	43.6%	43.6%	43.8%

Primary Endpoint: Mean Change in BCVA at Week 48

photon

DME



	Mean number of injections ^a	LS mean change from BL at Week 48 (MMRM)	Diff. in LS means vs. 2q8	2-sided 95% CI	1-sided test for non-inferiority at 4-letter margin
2q8	7.9	8.7			
8q12	6.0	8.1	-0.57	-2.26, 1.13	p < 0.0001
8q16	5.0	7.2	-1.44	-3.27, 0.39	p = 0.0031

Data shown in the figure represent LS mean values (censoring data post-ICE); FAS: 2q8 n=167; 8q12 n=328; 8q16 n=163 (at baseline).

LS mean values were generated using MMRM, with baseline BCVA as a covariate, treatment group (afibercept 2q8, 8q12, 8q16) and stratification variables (geographic region [Japan vs rest of the world], baseline CRT [<400 μm vs ≥400 μm], prior treatment for DME [yes vs no]) as fixed factors, and interaction terms for baseline and visit and for treatment and visit.


^aPatients completing Week 48: 2q8 n=157; 8q12 n=300; 8q16 n=156.

BL, baseline; ICE, intercurrent event; LS, least squares; MMRM, mixed model for repeated measures.


Large Majority of Aflibercept 8 mg Patients Maintained Randomized Intervals Through Week 48

photon


DME



8q12
(n=300)^b



8q16
(n=156)^b




All 8 mg
(n=456)^b

^aPatients whose dosing intervals were shortened based on DRM assessments at some point through Week 48.

^bPatients completing Week 48.


Exploratory Analysis of Patients Who Would Hypothetically Meet DRM Criteria for Additional Interval Shortening Through Week 48

1.7% of patients in the **8q12** group would have met DRM criteria for shortening to Q4




8q12
(n=300)^b

1.3% of patients in the **8q16** group would have met DRM criteria for shortening to Q4



8q16
(n=156)^b

1.5% of all 8 mg patients would have met DRM criteria for shortening to Q4



All 8 mg
(n=456)^b


In this exploratory analysis, eyes that were shortened to a Q8 dosing interval through Week 48 were further evaluated to determine if the study-specified DRM criteria would have been met for shortening to a Q4 interval. DRM criteria for interval shortening were defined as >10-letter loss in BCVA from Week 12 due to persistent or worsening DME and >50-μm increase in CRT from Week 12 (assessed using OCT data from the central reading center).

^aPatients whose dosing intervals were shortened based on DRM assessments at some point through Week 48.

^bPatients completing Week 48.

OCT, optical coherence tomography.

Mean Change in BCVA at Week 96



	Mean number of injections ^a	LS mean change from BL at Week 96 (MMRM)	Diff. in LS means vs. 2q8	2-sided 95% CI	1-sided test for non-inferiority at 4-letter margin
2q8	13.8	7.7			
8q12	9.5	8.2	+0.45	-1.55, 2.45	p < 0.0001 (nominal)
8q16	7.8	6.6	-1.11	-3.27, 1.05	p = 0.0044 (nominal)

Data shown in the figure represent LS mean values (censoring data post-ICE); FAS: 2q8 n=167; 8q12 n=328; 8q16 n=163 (at baseline).


LS mean values were generated using MMRM, with baseline BCVA as a covariate, treatment group (aflibercept 2q8, 8q12, 8q16) and stratification variables (geographic region [Japan vs rest of the world], baseline CRT [<400 μm vs ≥400 μm], prior treatment for DME [yes vs no]) as fixed factors, and interaction terms for baseline and visit and for treatment and visit.

^aPatients completing Week 96: 2q8 n=139; 8q12 n=256; 8q16 n=139.

BL, baseline; ICE, intercurrent event; LS, least squares; MMRM, mixed model for repeated measures.


Large Majority of Aflibercept 8 mg Patients Maintained Randomized Intervals Through Week 96

88% of patients in the 8q12 group maintained ≥ 12 -week intervals




Randomized to **8q12** at BL
(n=256)^b

84% of patients in the 8q16 group maintained ≥ 16 -week intervals



Randomized to **8q16** at BL
(n=139)^b




All 8 mg
(n=395)^b


^aPatients met DRM criteria for dosing interval shortening at some point through Week 96. ^bPatients completing Week 96. Values may not add up to 100% due to rounding.

Large Proportion of Patients Qualified for Interval Extension in Year 2^a

Last Completed




Last Assigned



^aDosing intervals were extended in Year 2 if patients had <5-letter loss in BCVA from Week 12 and CRT <300 μm (or <320 μm on Spectralis). ^bPatients were assigned to 24-week dosing intervals if they continued to meet extension criteria but there was not sufficient time to complete the interval within the 96-week study period. ^cPatients completing Week 96. Values may not add up to 100% due to rounding.


Dosing Interval for Majority of 8 mg Patients Was Maintained or Extended Through Week 96

8q12 (n=256)^a



43% of patients had a last assigned dosing interval of ≥ 20 weeks

8q16 (n=139)^a




47% of patients had a last assigned dosing interval of ≥ 20 weeks

■ Q8 ■ Q12 ■ Q16 ■ Q20 ■ Q24

^aPatients completing Week 96.

Mean Change in Central Retinal Thickness



LS mean values (censoring data post-ICE); FAS: 2q8 n=167; 8q12 n=328; 8q16 n=163 (at baseline).

LS mean values were generated using MMRM, with baseline CRT as a covariate, treatment group (aflibercept 2q8, 8q12, 8q16) and stratification variables (geographic region [Japan vs rest of the world], baseline CRT [$<400\ \mu\text{m}$ vs $\geq 400\ \mu\text{m}$], prior treatment for DME [yes vs no]) as fixed factors, and interaction terms for baseline and visit and for treatment and visit.

Safety Through Week 96

2q8

8q12

8q16

All 8 mg

DME

N (SAF)	167	328	163	491
Ocular safety				
Patients with ≥ 1 ocular AE (%) ^a	37.1%	43.9%	45.4%	44.4%
Patients with ≥ 1 IOI AE (%) ^a	1.2%	1.5%	0.6%	1.2%
Patients with IOP ≥ 35 mmHg pre- or post-injection (%) ^b	1.2%	0.6%	0	0.4%
Non-ocular safety				
APTC events (%) ^a	7.2%	6.7%	6.7%	6.7%
Hypertension events (%) ^a	16.2%	15.5%	20.9%	17.3%
Non-ocular SAEs (%) ^a	25.1%	22.9%	23.9%	23.2%
Deaths (%) ^c	5.4%	5.5%	3.1%	4.7%

- Ocular AEs occurring in $\geq 5\%$ of patients in any treatment group were cataract, vitreous floaters, and conjunctival hemorrhage
- No cases of ischemic optic neuropathy, retinal vasculitis, or occlusive retinitis were reported through Week 96
- Mean changes from baseline in pre-dose IOP did not exceed ± 1 mmHg at any timepoint through Week 96 in any treatment group

^aTreatment emergent. ^bIOP was measured in the study eye. ^cAll events.

AE, adverse event; APTC, Anti-Platelet Trialists' Collaboration; IOI, intraocular inflammation; SAE, serious adverse event.


PHOTON: 96-week Results



DME


- 8q12 and 8q16 groups achieved similar BCVA gains compared to 2q8 at Week 96, with up to 6 fewer injections
- Through Week 96, 88% of 8q12 patients and 84% of 8q16 patients maintained ≥ 12 - and ≥ 16 -week dosing intervals, respectively
 - At Week 96, 43% of 8q12 patients and 47% of 8q16 patients had a last assigned dosing interval of ≥ 20 weeks
- Safety of aflibercept 8 mg was comparable to that of aflibercept 2 mg over 96 weeks

Mean Change From Baseline in BCVA^a



	LS mean change from BL at Week 96 (MMRM)	Diff. in LS means vs. 2q8	2-sided 95% CI	1-sided test for NI at 4-letter margin
2q8	7.7			
8q12	8.2	+0.45	-1.55, 2.45	p < 0.0001 (nominal)
8q16	6.6	-1.11	-3.27, 1.05	p = 0.0044 (nominal)

Last Assigned Dosing Interval at Week 96



^aLS mean values (censoring data post-ICE); FAS: 2q8 n=167; 8q12 n=328; 8q16 n=163 (at baseline). LS mean values were generated using MMRM, with baseline BCVA as a covariate, treatment group (aflibercept 2q8, 8q12, 8q16) and stratification variables (geographic region [Japan vs rest of the world], baseline CRT [$<400 \mu\text{m}$ vs $\geq 400 \mu\text{m}$], prior treatment for DME [yes vs no]) as fixed factors, and interaction terms for baseline and visit and for treatment and visit. ^bPatients were assigned to 24-week dosing intervals if they continued to meet extension criteria but there was not sufficient time to complete the interval within the 96-week study period. ^cPatients completing Week 96.

Thank you to all co-investigators and patients who participated in the PHOTON trial

photon

DME

