



photon

**Week 48 Outcomes in Aflibercept 8 mg- and 2 mg-treated Patients
by Prior DME Treatment Status:
A Subgroup Analysis of the Phase 2/3 PHOTON Trial**

Nitish Mehta MD, on behalf of the PHOTON study investigators

Department of Ophthalmology, NYU Langone Health, New York, New York

Presented at Association for Research in Vision and Ophthalmology, May 5–9, 2024

Background

- Aflibercept 8 mg is a novel intravitreal formulation that delivers a 4-times higher molar dose than aflibercept 2 mg, potentially extending VEGF suppression over a longer period of time
- In the PHOTON trial, aflibercept 8 mg demonstrated non-inferior BCVA gains with extended dosing intervals versus aflibercept 2 mg in patients with DME, with no new safety signals through Week 48¹
 - Given that approximately 44% of patients in PHOTON received prior treatment for DME,^a there is an opportunity to assess treatment outcomes in patients with prior DME treatment

This subgroup analysis of the PHOTON trial evaluated visual and anatomic outcomes in patients by prior DME treatment status

^aPrior DME treatment status was categorized as yes/no in the electronic data capture record. Previous treatments for DME were laser, intravitreal anti-VEGF therapy, and corticosteroids.

BCVA, best-corrected visual acuity; DME, diabetic macular edema; VEGF, vascular endothelial growth factor.

1. Brown DM. *Lancet*. 2024:S0140–6736(23)02577–1. Online ahead of print.

PHOTON Study Design

photon

DME

Multi-center, randomized, double-masked study in patients with DME^a
Randomized 1 (2q8) : 2 (8q12) : 1 (8q16)

2q8

Aflibercept 2q8
after 5 initial monthly injections
n=167

8q12

Aflibercept 8q12
after 3 initial monthly injections
n=328

8q16

Aflibercept 8q16
after 3 initial monthly injections
n=163

Primary endpoint at Week 48:
Mean change in BCVA (non-inferiority)

Key secondary endpoint:
Proportion of patients with ≥ 2 -step improvement in DRSS at Week 48



End of study at Week 96
with optional 1-year extension through Week 156

^aTreatment naïve and previously treated.
DRSS, Diabetic Retinopathy Severity Scale.

PHOTON: Dose Regimen Modifications in Year 1

	Day 1	Week 4	Week 8	Week 12	Week 16	Week 20	Week 24	Week 28	Week 32	Week 36	Week 40	Week 44	Week 48
2q8	X	X	X	X	X	o	X	o	X	o	X	o	X
8q12	X	X	X	o	o	X	o	o	X	o	o	X	o
8q16	X	X	X	o	o	o	X	o	o	o	X	o	o

DRM Criteria for Shortening Dosing Interval^a

- >10-letter loss in BCVA due to persistent or worsening DME
- AND
- >50-micron increase in CRT

DRM in Year 1

Intervals can only be **shortened**

Multiple opportunities to shorten interval

Minimum interval for all patients was **Q8**

Week 16 and 20: Patients on **8q12** and **8q16** meeting DRM criteria shortened to Q8

Week 24: Patients on **8q16** meeting DRM criteria shortened to Q12

Week 32 and 44 for **8q12** and Week 40 for **8q16**: Treatment interval shortened by 4 weeks for patients meeting DRM criteria

Stippled boxes = initial treatment phase; X = active injection; o = sham injections. Note: Figure does not reflect all dosing options once a patient's dosing interval is shortened.

^aAll assessments compared to Week 12.

CRT, central retinal thickness; DRM, dose regimen modification; Q8, every 8 weeks; Q12, every 12 weeks.

Baseline Demographics

With Prior DME Treatment

Without Prior DME Treatment

Age, years
Female, %
Race, %
White
Asian
Black or African American
American Indian or Alaskan Native
Other
Not reported
Hispanic or Latino, %
Duration of diabetes, years

2q8 (n=74)	8q12 (n=143)	8q16 (n=71)	2q8 (n=93)	8q12 (n=185)	8q16 (n=92)
64.4 (8.9)	62.8 (11.0)	63.0 (8.4)	62.0 (10.4)	61.6 (11.3)	60.9 (10.3)
45.9	39.2	40.8	44.1	33.5	38.0
64.9	69.9	77.5	68.8	70.8	79.3
21.6	19.6	18.3	15.1	10.8	10.9
9.5	7.0	4.2	11.8	13.5	6.5
0.0	0.7	0.0	0.0	0.5	0.0
2.7	1.4	0.0	2.2	2.2	1.1
1.4	1.4	0.0	2.2	1.1	2.2
18.9	17.5	22.5	18.3	15.7	19.6
16.7 (10.6)	16.0 (9.4)	16.6 (9.7)	15.3 (9.6)	14.5 (10.3)	15.0 (11.4)


Baseline Ocular Characteristics

	With Prior DME Treatment			Without Prior DME Treatment		
	2q8 (n=74)	8q12 (n=143)	8q16 (n=71)	2q8 (n=93)	8q12 (n=185)	8q16 (n=92)
BCVA, ETDRS letters	62.1 (10.9)	62.3 (10.5)	58.6 (11.9)	61.0 (11.5)	64.7 (9.7)	63.7 (11.2)
Snellen equivalent, %						
20/32 (>73 to 78 letters)	14.9	16.8	5.6	9.7	18.4	20.7
20/40 or worse ($\leq$ 73 letters)	85.1	83.2	94.4	90.3	81.1	79.3
CRT, μm	472.7 (162.3)	455.7 (124.0)	460.6 (109.3)	444.9 (127.1)	444.1 (130.1)	460.1 (124.7)
DRSS categories, %						
Better than or equal to level 43	70.3	66.4	67.6	57.0	55.1	64.1
Level 47 or worse	25.7	28.0	23.9	36.6	39.5	31.5
Missing/ungradable	4.1	5.6	8.5	6.5	5.4	4.3


Mean Change in BCVA Through Week 48

With Prior DME Treatment

DME




Without Prior DME Treatment




Proportion of Patients With ≥ 2 -step DRSS Improvement From Baseline at Week 48

With Prior DME Treatment




Without Prior DME Treatment




Proportion of Patients With Baseline DRSS 47 or Worse and ≥ 2 -step DRSS Improvement From Baseline at Week 48

With Prior DME Treatment




Without Prior DME Treatment




Mean Change in CRT Through Week 48

With Prior DME Treatment

DME




Without Prior DME Treatment




Proportion of Patients Who Maintained Their Randomized Intervals Through Week 48

With Prior DME Treatment



Without Prior DME Treatment




Values may not add up to 100% due to rounding.

^aPatients whose dosing intervals were shortened based on DRM assessments at some point through Week 48.


^bPatients completing Week 48.

Conclusions


Proportion of Patients With ≥ 2 -step DRSS Improvement From Baseline at Week 48



Mean Change in BCVA Through Week 48



Proportion of Patients Who Maintained Their Randomized Intervals Through Week 48



- Outcomes were generally comparable across treatment groups within subgroups of patients with or without prior DME treatment
- BCVA gains and proportions of patients with ≥ 2 -step improvement in DRSS score at Week 48 trended numerically higher across all treatment groups in patients without versus with prior DME treatment
- Similar proportions of 8q12 and 8q16 patients maintained ≥ 12 -week dosing through Week 48 irrespective of prior DME treatment status