

# **Impact of Baseline Best-corrected Visual Acuity on Visual and Anatomic Outcomes in Patients With Retinal Vein Occlusion: QUASAR Post Hoc Analysis**

**Roger Goldberg, MD, on behalf of the QUASAR study investigators**

*Bay Area Retina Associates, Walnut Creek, California*

## Disclosures

- Roger Goldberg received consulting and/or speaking fees from AbbVie, Allergan, Annexon, Apellis, Biogen, Boehringer Ingelheim, Carl Zeiss Meditech, Genentech/Roche, and Regeneron Pharmaceuticals, Inc.; received research/grant support from Allergan/AbbVie, Aerie, Alexion, Annexon, Apellis, Boehringer Ingelheim, Carl Zeiss Meditec, Eyepoint, Genentech/Roche, Janssen, Novo Nordisk, Ocuphire, and Unity Bio; and holding equity in Emmetrope Ophthalmics
- The QUASAR trial (NCT05850520) was sponsored by Bayer AG (Leverkusen, Germany). This analysis was funded by Regeneron Pharmaceuticals, Inc. (Tarrytown, New York). The sponsor participated in the design and conduct of this analysis, interpretation of the data, and preparation of this presentation
- Study disclosures: This study included research conducted on human patients. Institutional Review Board/Institutional Ethics Committee approval was obtained prior to study initiation
- The data in this presentation were originally presented at the ARVO Annual Meeting, May 3–7, 2026, Denver, CO, USA
- Medical writing support was provided by MEDiSTRAVA (London, UK), funded by Regeneron Pharmaceuticals, Inc., according to Good Publication Practice (GPP) guidance (*Ann Intern Med.* 2022;175:1298-1304)

## Background and Objective

- In the QUASAR trial, patients treated with aflibercept 8 mg achieved noninferior BCVA gains and comparable reductions in CRT with fewer injections versus aflibercept 2 mg through Week 36, overall and across RVO subtypes (BRVO or CRVO/HRVO)<sup>1</sup>
- Understanding factors that may impact visual and anatomic outcomes can help guide treatment decisions and manage physician and patient expectations

**This post hoc analysis evaluated the impact of baseline BCVA on visual and anatomic outcomes in patients with macular edema secondary to BRVO or CRVO/HRVO after treatment with aflibercept 8 mg versus 2 mg in QUASAR**

# QUASAR: Study Design

A multicenter, randomized, double-masked, phase 3 study in patients with treatment-naïve macular edema secondary to RVO  
Randomized at baseline 1 (2q4) : 1 (8q8/3) : 1 (8q8/5)

**2q4**  
Aflibercept 2 mg initiated with 9 monthly injections, followed by T&E<sup>a</sup>  
n=301

**8q8/3**  
Aflibercept 8 mg initiated with 3 monthly injections, followed by Q8 and T&E<sup>a</sup>  
n=293

**8q8/5**  
Aflibercept 8 mg initiated with 5 monthly injections, followed by Q8 and T&E<sup>a</sup>  
n=298

**Primary endpoint**  
Mean change in BCVA (noninferiority)

|       | Day 1 | W4 | W8 | W12 | W16 | W20            | W24 | W28            | W32 | W36            |
|-------|-------|----|----|-----|-----|----------------|-----|----------------|-----|----------------|
| 2q4   | X     | X  | X  | X   | X   | X              | X   | X              | X   | T&E            |
| 8q8/3 | X     | X  | X  | o   | X   | o <sup>b</sup> | X   | o <sup>c</sup> | X   | T&E            |
| 8q8/5 | X     | X  | X  | X   | X   | o              | X   | o <sup>c</sup> | X   | o <sup>d</sup> |

## DRM for interval shortening

Dosing interval shortened by 4 weeks beginning at Week 16 for 8q8/3, Week 24 for 8q8/5, or Week 40 for 2q4 if both the following criteria were met at a dosing visit:

- BCVA loss of >5 letters from the reference visit<sup>d</sup>, AND
- >50-µm increase in CRT from the reference visit<sup>d</sup>

## DRM for interval extension

Dosing interval extended by 4 weeks starting at Week 32 for 8q8/3 and 2q4 and at Week 40 for 8q8/5 if both the following criteria were met at a dosing visit:

- BCVA loss of <5 letters from the reference visit<sup>d</sup>, AND
- CRT <320 µm on Heidelberg (or <300 µm on Cirrus or Topcon SD-OCT)

The primary efficacy endpoint was change from baseline in BCVA at Week 36, with a non-inferiority margin of 4 letters. Stippled boxes = initial treatment phase; X = active injection; o = sham injection. Note: Table does not reflect all dosing options once a patient's dosing interval is shortened. <sup>a</sup>With opportunity for extension per DRM. <sup>b</sup>Active injection for patients meeting DRM criteria at Week 16. <sup>c</sup>Active injection for patients meeting DRM criteria from Week 16 for 8q8/3 or Week 24 for 8q8/5. <sup>d</sup>Reference is Week 12 visit for 8q8/3 and Week 20 visit for 8q8/5 and 2q4 (denoted by green boxes on table). 2q4, aflibercept 2 mg every 4 weeks; 8q8/3, aflibercept 8 mg every 8 weeks after 3 initial injections at 4-week intervals; 8q8/5, aflibercept 8 mg every 8 weeks after 5 initial injections at 4-week intervals; DRM, dose regimen modification; Q8, every 8 weeks; SD-OCT, spectral-domain optical coherence tomography; T&E, treat and extend; W, week.

## Methods

- A tertile analysis by baseline BCVA was conducted post hoc to evaluate visual and anatomic outcomes through Week 36 following treatment with aflibercept 8 mg or 2 mg in patients with BRVO or CRVO/HRVO from the QUASAR trial
- Patients were grouped by baseline BCVA as follows:

### BRVO

**T1:  $\leq 55$  ETDRS letters**

**T2:  $>55$  to  $\leq 65$  ETDRS letters**

**T3:  $>65$  ETDRS letters**

### CRVO/HRVO

**T1:  $\leq 47$  ETDRS letters**

**T2:  $>47$  to  $\leq 59$  ETDRS letters**

**T3:  $>59$  ETDRS letters**

- Data were summarized descriptively

**Visual and Anatomic Outcomes  
by Baseline BCVA Tertiles  
in Patients With BRVO**

# Baseline Demographics and Disease Characteristics by Baseline BCVA

BRVO

|                                                                                  | T1: ≤55 ETDRS letters |                  |                  | T2: >55 to ≤65 ETDRS letters |                  |                  | T3: >65 ETDRS letters |                  |                  |
|----------------------------------------------------------------------------------|-----------------------|------------------|------------------|------------------------------|------------------|------------------|-----------------------|------------------|------------------|
|                                                                                  | 2q4<br>(n=54)         | 8q8/3<br>(n=50)  | 8q8/5<br>(n=53)  | 2q4<br>(n=44)                | 8q8/3<br>(n=48)  | 8q8/5<br>(n=56)  | 2q4<br>(n=51)         | 8q8/3<br>(n=61)  | 8q8/5<br>(n=50)  |
| <b>Age, mean (SD), years</b>                                                     | 66.9<br>(10.6)        | 67.0<br>(10.8)   | 67.6<br>(11.0)   | 65.9<br>(9.6)                | 65.7<br>(10.4)   | 65.2<br>(12.0)   | 61.9<br>(11.4)        | 63.3<br>(10.8)   | 61.9<br>(11.0)   |
| <b>Female, n (%)</b>                                                             | 28 (51.9)             | 28 (56.0)        | 34 (64.2)        | 25 (56.8)                    | 26 (54.2)        | 30 (53.6)        | 25 (49.0)             | 29 (47.5)        | 23 (46.0)        |
| <b>Race, n (%)</b>                                                               |                       |                  |                  |                              |                  |                  |                       |                  |                  |
| White                                                                            | 30 (55.6)             | 26 (52.0)        | 30 (56.6)        | 20 (45.5)                    | 22 (45.8)        | 23 (41.1)        | 31 (60.8)             | 39 (63.9)        | 32 (64.0)        |
| Asian                                                                            | 24 (44.4)             | 19 (38.0)        | 20 (37.7)        | 19 (43.2)                    | 20 (41.7)        | 30 (53.6)        | 18 (35.3)             | 18 (29.5)        | 13 (26.0)        |
| Other <sup>a</sup>                                                               | 0                     | 0                | 1 (1.9)          | 1 (2.3)                      | 1 (2.1)          | 1 (1.8)          | 0                     | 1 (1.6)          | 3 (6.0)          |
| Not reported                                                                     | 0                     | 5 (10.0)         | 2 (3.8)          | 4 (9.1)                      | 5 (10.4)         | 2 (3.6)          | 2 (3.9)               | 3 (4.9)          | 2 (4.0)          |
| <b>Hispanic or Latino, n (%)</b>                                                 | 5 (9.3)               | 4 (8.0)          | 2 (3.8)          | 2 (4.5)                      | 5 (10.4)         | 3 (5.4)          | 4 (7.8)               | 2 (3.3)          | 3 (6.0)          |
| <b>History of hypertension, n (%)</b>                                            | 31 (57.4)             | 32 (64.0)        | 32 (60.4)        | 30 (68.2)                    | 32 (66.7)        | 39 (69.6)        | 27 (52.9)             | 40 (65.6)        | 32 (64.0)        |
| <b>BCVA, mean (SD), ETDRS letters</b>                                            | 42.6 (9.9)            | 44.5 (7.9)       | 45.1 (8.1)       | 61.1 (3.0)                   | 59.9 (3.0)       | 61.0 (3.2)       | 69.6 (2.1)            | 69.2 (2.2)       | 70.0 (2.3)       |
| <b>CRT, mean (SD), μm</b>                                                        | 647.1<br>(183.6)      | 623.1<br>(171.8) | 603.9<br>(191.6) | 526.6<br>(136.3)             | 556.1<br>(157.1) | 518.1<br>(143.9) | 475.0<br>(131.2)      | 482.2<br>(153.9) | 498.7<br>(122.8) |
| <b>Total area of macular ischemia,<sup>b</sup><br/>mean (SD), mm<sup>2</sup></b> | 6.9 (3.9)             | 6.8 (3.4)        | 6.7 (3.5)        | 6.3 (3.6)                    | 5.8 (3.7)        | 5.6 (3.9)        | 4.8 (3.6)             | 4.5 (3.7)        | 5.7 (4.5)        |

FAS.


<sup>a</sup>Includes patients who were Black or African American, Native Hawaiian or other Pacific Islander, or multiracial. <sup>b</sup>Based on FA, not considering the foveal avascular zone.

FA, fluorescein angiography; FAS, full analysis set.


# Mean BCVA Through Week 36 by Baseline BCVA

BRVO


T1: ≤55 ETDRS letters



T2: >55 to ≤65 ETDRS letters



T3: >65 ETDRS letters



Mean number of injections<sup>a</sup>

Mean change in BCVA at Week 36, ETDRS letters

|              |     |       |
|--------------|-----|-------|
| 2q4 (n=54)   | 8.9 | +27.2 |
| 8q8/3 (n=50) | 6.0 | +24.8 |
| 8q8/5 (n=53) | 6.9 | +25.8 |

Mean number of injections<sup>a</sup>

Mean change in BCVA at Week 36, ETDRS letters

|              |     |       |
|--------------|-----|-------|
| 2q4 (n=44)   | 8.9 | +16.7 |
| 8q8/3 (n=48) | 6.1 | +17.3 |
| 8q8/5 (n=56) | 7.0 | +18.9 |

Mean number of injections<sup>a</sup>

Mean change in BCVA at Week 36, ETDRS letters

|              |     |       |
|--------------|-----|-------|
| 2q4 (n=51)   | 8.7 | +13.6 |
| 8q8/3 (n=61) | 6.1 | +12.1 |
| 8q8/5 (n=50) | 6.9 | +13.5 |

In BRVO, aflibercept 8 mg achieved robust improvements in BCVA through Week 36 with fewer injections than aflibercept 2 mg across baseline BCVA tertiles


FAS, observed cases excluding intercurrent events.

<sup>a</sup>Patients who completed the Week 36 visit.


# Mean CRT Through Week 36 by Baseline BCVA

BRVO


T1: ≤55 ETDRS letters



T2: >55 to ≤65 ETDRS letters



T3: >65 ETDRS letters



Mean number of injections<sup>a</sup>

Mean change in CRT at Week 36, μm

2q4 (n=54) 8.9 -400.4

8q8/3 (n=50) 6.0 -361.1

8q8/5 (n=53) 6.9 -355.3

Mean number of injections<sup>a</sup>

Mean change in CRT at Week 36, μm

2q4 (n=44) 8.9 -279.0

8q8/3 (n=48) 6.1 -301.0

8q8/5 (n=56) 7.0 -265.7

Mean number of injections<sup>a</sup>

Mean change in CRT at Week 36, μm

2q4 (n=51) 8.7 -200.2

8q8/3 (n=61) 6.1 -213.3

8q8/5 (n=50) 6.9 -232.4


In BRVO, aflibercept 8 mg achieved robust CRT reductions through Week 36 with fewer injections than aflibercept 2 mg across baseline BCVA tertiles

FAS, observed cases excluding intercurrent events.

<sup>a</sup>Patients who completed the Week 36 visit.

# Last Assigned Dosing Interval at Week 36 by Baseline BCVA

BRVO



At Week 36, most patients treated with aflibercept 8 mg achieved a last assigned dosing interval of ≥Q8, and fewer patients required Q4 dosing versus 2 mg across baseline BCVA tertiles

Patients who completed the Week 36 visit. Values may not add up to 100% because of rounding.

<sup>a</sup>Per study design, dosing interval extension was not possible in the 8q8/5 group until Week 40.

Q4, every 4 weeks; Q12, every 12 weeks.

**Visual and Anatomic Outcomes  
by Baseline BCVA Tertiles  
in Patients With CRVO/HRVO**

# Baseline Demographics and Disease Characteristics by Baseline BCVA

CRVO/  
HRVO

|                                                                                  | T1: ≤47 ETDRS letters |                  |                  |  | T2: >47 to ≤59 ETDRS letters |                  |                  |  | T3: >59 ETDRS letters |                  |                  |
|----------------------------------------------------------------------------------|-----------------------|------------------|------------------|--|------------------------------|------------------|------------------|--|-----------------------|------------------|------------------|
|                                                                                  | 2q4<br>(n=54)         | 8q8/3<br>(n=45)  | 8q8/5<br>(n=42)  |  | 2q4<br>(n=44)                | 8q8/3<br>(n=43)  | 8q8/5<br>(n=53)  |  | 2q4<br>(n=54)         | 8q8/3<br>(n=46)  | 8q8/5<br>(n=44)  |
| <b>Age, mean (SD), years</b>                                                     | 68.4<br>(12.9)        | 69.1<br>(12.5)   | 69.3<br>(10.7)   |  | 66.4<br>(10.5)               | 65.2<br>(13.1)   | 67.4<br>(12.8)   |  | 66.0<br>(13.7)        | 65.6<br>(11.6)   | 63.5<br>(9.8)    |
| <b>Female, n (%)</b>                                                             | 26 (48.1)             | 23 (51.1)        | 20 (47.6)        |  | 20 (45.5)                    | 15 (34.9)        | 25 (47.2)        |  | 20 (37.0)             | 15 (32.6)        | 14 (31.8)        |
| <b>Race, n (%)</b>                                                               |                       |                  |                  |  |                              |                  |                  |  |                       |                  |                  |
| White                                                                            | 39 (72.2)             | 27 (60.0)        | 31 (73.8)        |  | 25 (56.8)                    | 26 (60.5)        | 32 (60.4)        |  | 33 (61.1)             | 33 (71.7)        | 29 (65.9)        |
| Asian                                                                            | 12 (22.2)             | 10 (22.2)        | 7 (16.7)         |  | 15 (34.1)                    | 14 (32.6)        | 17 (32.1)        |  | 13 (24.1)             | 10 (21.7)        | 10 (22.7)        |
| Other <sup>a</sup>                                                               | 2 (3.7)               | 3 (6.7)          | 2 (4.8)          |  | 2 (4.5)                      | 1 (2.3)          | 2 (3.8)          |  | 4 (7.4)               | 1 (2.2)          | 4 (9.1)          |
| Not reported                                                                     | 1 (1.9)               | 5 (11.1)         | 2 (4.8)          |  | 2 (4.5)                      | 2 (4.7)          | 2 (3.8)          |  | 4 (7.4)               | 2 (4.3)          | 1 (2.3)          |
| <b>Hispanic or Latino, n (%)</b>                                                 | 2 (3.7)               | 6 (13.3)         | 3 (7.1)          |  | 6 (13.6)                     | 4 (9.3)          | 1 (1.9)          |  | 3 (5.6)               | 4 (8.7)          | 2 (4.5)          |
| <b>History of hypertension, n (%)</b>                                            | 35 (64.8)             | 32 (71.1)        | 33 (78.6)        |  | 28 (63.6)                    | 25 (58.1)        | 27 (50.9)        |  | 36 (66.7)             | 31 (67.4)        | 33 (75.0)        |
| <b>BCVA, mean (SD), ETDRS letters</b>                                            | 33.7 (7.9)            | 33.7 (8.1)       | 32.7 (7.0)       |  | 54.3 (3.1)                   | 53.1 (3.1)       | 54.3 (3.5)       |  | 65.6 (4.0)            | 66.7 (4.2)       | 67.1 (3.8)       |
| <b>CRT, mean (SD), μm</b>                                                        | 914.4<br>(287.7)      | 895.4<br>(278.7) | 837.3<br>(251.4) |  | 730.5<br>(195.8)             | 700.4<br>(213.5) | 679.5<br>(198.2) |  | 598.9<br>(166.3)      | 560.6<br>(142.2) | 554.8<br>(183.0) |
| <b>Total area of macular ischemia,<sup>b</sup><br/>mean (SD), mm<sup>2</sup></b> | 4.0 (7.3)             | 6.1 (7.4)        | 5.0 (7.7)        |  | 2.3 (4.3)                    | 2.9 (5.3)        | 2.7 (4.4)        |  | 1.3 (2.6)             | 2.2 (3.4)        | 1.6 (3.3)        |


FAS.

<sup>a</sup>Includes patients who were Black or African American, American Indian or Alaska Native, or multiracial. <sup>b</sup>Based on FA, not considering the foveal avascular zone.


# Mean BCVA Through Week 36 by Baseline BCVA

CRVO/  
HRVO


**T1: ≤47 ETDRS letters**



**T2: >47 to ≤59 ETDRS letters**



**T3: >59 ETDRS letters**



Mean number  
of injections<sup>a</sup>

Mean change in  
BCVA at Week 36,  
ETDRS letters

|              |     |       |
|--------------|-----|-------|
| 2q4 (n=54)   | 8.7 | +19.6 |
| 8q8/3 (n=45) | 6.1 | +19.9 |
| 8q8/5 (n=42) | 6.9 | +25.9 |

Mean number  
of injections<sup>a</sup>

Mean change in  
BCVA at Week 36,  
ETDRS letters

|              |     |       |
|--------------|-----|-------|
| 2q4 (n=44)   | 8.6 | +17.9 |
| 8q8/3 (n=43) | 6.1 | +17.6 |
| 8q8/5 (n=53) | 7.0 | +17.8 |

Mean number  
of injections<sup>a</sup>

Mean change in  
BCVA at Week 36,  
ETDRS letters

|              |     |       |
|--------------|-----|-------|
| 2q4 (n=54)   | 8.9 | +11.6 |
| 8q8/3 (n=46) | 6.1 | +12.5 |
| 8q8/5 (n=44) | 6.9 | +12.9 |

**In CRVO/HRVO, aflibercept 8 mg achieved robust improvements in BCVA through Week 36 with fewer injections than aflibercept 2 mg across baseline BCVA tertiles**


FAS, observed cases excluding intercurrent events.

<sup>a</sup>Patients who completed the Week 36 visit.


# Mean CRT Through Week 36 by Baseline BCVA

CRVO/  
HRVO


**T1: ≤47 ETDRS letters**



**T2: >47 to ≤59 ETDRS letters**



**T3: >59 ETDRS letters**



Mean number  
of injections<sup>a</sup>

Mean change in  
CRT at Week 36,  
μm

|              |     |        |
|--------------|-----|--------|
| 2q4 (n=54)   | 8.7 | -688.0 |
| 8q8/3 (n=45) | 6.1 | -665.2 |
| 8q8/5 (n=42) | 6.9 | -593.2 |

Mean number  
of injections<sup>a</sup>

Mean change in  
CRT at Week 36,  
μm

|              |     |        |
|--------------|-----|--------|
| 2q4 (n=44)   | 8.6 | -468.6 |
| 8q8/3 (n=43) | 6.1 | -449.0 |
| 8q8/5 (n=53) | 7.0 | -440.3 |

Mean number  
of injections<sup>a</sup>

Mean change in  
CRT at Week 36,  
μm

|              |     |        |
|--------------|-----|--------|
| 2q4 (n=54)   | 8.9 | -345.4 |
| 8q8/3 (n=46) | 6.1 | -292.1 |
| 8q8/5 (n=44) | 6.9 | -265.3 |


**In CRVO/HRVO, aflibercept 8 mg achieved robust CRT reductions through Week 36 with fewer injections than aflibercept 2 mg across baseline BCVA tertiles**

FAS, observed cases excluding intercurrent events.

<sup>a</sup>Patients who completed the Week 36 visit.

# Last Assigned Dosing Interval at Week 36 by Baseline BCVA

CRVO/  
HRVO



At Week 36, most patients treated with aflibercept 8 mg achieved a last assigned dosing interval of ≥Q8, and fewer patients required Q4 dosing versus 2 mg across baseline BCVA tertiles

Patients who completed the Week 36 visit. Values may not add up to 100% because of rounding.

<sup>a</sup>Per study design, dosing interval extension was not possible in the 8q8/5 group until Week 40.

Q4, every 4 weeks; Q12, every 12 weeks.

## Conclusions

- In patients with BRVO or CRVO/HRVO, aflibercept 8 mg achieved robust gains in BCVA and reductions in CRT with fewer injections versus aflibercept 2 mg through Week 36, regardless of baseline BCVA
- Patients with lower BCVA achieved greater visual gains, however their final BCVA was lower than those who started with better vision
- The majority of patients with BRVO or CRVO/HRVO treated with aflibercept 8 mg achieved extended dosing intervals regardless of baseline BCVA, with 93%-95% and 87%-98%, respectively, having a last assigned interval of  $\geq$ Q8 at Week 36
  - Substantially fewer patients treated with aflibercept 8 mg had a last assigned dosing interval of Q4 versus those treated with aflibercept 2 mg at Week 36